

# A refined analysis of the remarkable Bp star HR 6000 ★

F. Castelli<sup>1</sup> and S. Hubrig<sup>2</sup>

<sup>1</sup> Istituto Nazionale di Astrofisica– Osservatorio Astronomico di Trieste, Via Tiepolo 11, I-34131 Trieste, Italy  
e-mail: castelli@oats.inaf.it

<sup>2</sup> European Southern Observatory, Casilla 19001, Santiago 19, Chile  
e-mail: shubrig@eso.org

## ABSTRACT

**Aims.** UVES spectra of the very young ( $\sim 10^7$  years) peculiar B-type star HR 6000 were analyzed in the near-UV and visual spectral regions (3050–9460 Å) with the aim to extend to other spectral ranges the study made previously in the UV using IUE spectra.

**Methods.** Stellar parameters  $T_{\text{eff}}=12850$  K,  $\log g=4.10$ , and  $\xi=0$  km s<sup>-1</sup>, as determined from H $\beta$ , H $\gamma$ , H $\delta$  Balmer profiles and from the Fe I, Fe II ionization equilibrium, were used to compute an individual abundances ATLAS12 model. We identified spectral peculiarities and obtained final stellar abundances by comparing observed and computed equivalent widths and line profiles.

**Results.** The adopted model fails to reproduce the (b-y) and c color indices. The spectral analysis has revealed: the presence of emission lines for Mn II, Cr II, and Fe II; isotopic anomalies for Hg, Ca; the presence of interstellar lines of Na I at  $\lambda\lambda$  3302.3, 3302.9, 5890, 5896 Å, and of K I at 7665, 7699 Å; the presence of a huge quantity of unidentified lines, which we presume to be mostly due to Fe II transitions owing to the large Fe overabundance amounting to [+0.7]. The main chemical peculiarities are an extreme overabundance of Xe, followed by those of Hg, P, Y, Mn, Fe, Be, and Ti. The most underabundant element is Si, followed by C, N, Al, S, Mg, V, Sr, Co, Cl, Sc, and Ni. The silicon underabundance [-2.9] is the lowest value for Si ever observed in any HgMn star. The observed lines of He I can not be reproduced by a single value of the He abundance, but they require values ranging from [-0.8] to [-1.6]. Furthermore, when the observed and computed wings of He I lines are fitted, the observed line cores are much weaker than the computed ones. From the present analysis we infer the presence of vertical abundance stratification for He, Mn, and possibly also P.

**Key words.** stars:abundances - line:identification-atomic data-stars:atmospheres-stars:chemically peculiar- stars:individual:HR 6000 (Bp)

## 1. Introduction

HR 6000 (HD 144667) is one of the most remarkable chemically peculiar (CP) stars. It does not fit any of the CP subclasses, but it seems to combine abundance anomalies from a variety of Bp sub-types. It forms with the star HR 5999 (HD 144668) the common proper motion visual binary system  $\Delta$ 199 or Dunlop 199 (Bessell & Eggen, 1972). The angular separation between the HR 6000, which is the brighter more massive component, and the secondary component, the well-known Herbig Ae star HR 5999, is about 45".

The main interest to study the chemical composition of HR 6000 comes from the generally poor understanding of the occurrence of abundance anomalies in such a young object with an estimated age of the order of  $10^7$  years. In fact, the  $\Delta$  199 double system is located close to the center of the Lupus 3 molecular cloud which is populated by numerous T Tauri stars. This has led to an assumption that this system has the same age as the cloud, which is estimated to be  $(9.1 \pm 2.1) \times 10^6$  years (James et al., 2006). However, after the Hipparcos mission, the membership of HR 5999 and HR 6000 to the cloud became rather questionable due to the uncertainties in the distance determination for the Lupus cloud. The distance of HR 5999 and HR 6000 measured by Hipparcos are  $208 \pm 38$  pc and  $240 \pm 48$  pc, respectively, while the Lupus cloud distance is  $150 \pm 10$  pc according

to Crawford (2000). Comerón et al. (2003) assigned a distance of about 200 pc to the Lupus cloud, but this determination was made by assuming a priori that  $\Delta$ 199 belongs to the cloud.

A ROSAT survey of Herbig Ae/Be stars presented by Zinnecker & Preibisch (1994) has detected strong X-ray emission coming from the direction of HR 6000. To explain the X-ray origin, a possible T Tauri companion for HR 6000 was postulated by van den Ancker et al. (1996) on the basis of an infrared excess in the energy distribution relative to predictions made for an effective temperature of  $T_{\text{eff}}=14000$  K. Siebenmorgen et al. (2000) fitted very short spectral regions of observed flux in the mid IR to a black body of 13000 K and noticed that the observed spectrum was featureless.

Neither spectrum nor radial velocity variability was found for HR 6000 by Andersen & Jaschek (1984), who studied optical spectra taken at different epochs. However, van den Ancker et al. (1996) discovered long-term variations in the u, v, and b Strömgren magnitudes with approximate amplitude of  $0^m.03 \pm 0^m.01$  in u,  $0^m.02 \pm 0^m.01$  in v, and  $0^m.01$  in b. No variations in y were found. Kurtz & Marang (1995) discovered variations of 0.008 mag in V with a period near to 2 d and suggested that this period could be possibly caused by rotational variation in a spotted magnetic star. This would imply a pole on orientation for HR 6000 as they measured  $v \sin i \leq 5$  km s<sup>-1</sup>.

Catanzaro et al. (2004) were the first ones who provided abundances from the optical range. Previously, optical spectra were studied only by Andersen & Jaschek (1984) and Andersen et al. (1984) who identified the spectra from 3323 Å

Send offprint requests to: F. Castelli

\* Based on observations collected at the European Southern Observatory, Paranal, Chile (ESO programme 076.D-0169(A)).

to 5317 Å and discussed the abundances on the basis of the line intensities. It is remarkable that the stellar parameters derived by Catanzaro et al. (2004) from the Balmer profiles ( $T_{\text{eff}}=12950\pm50$  K,  $\log g=4.05\pm0.01$ ) are 1000 K lower than those previously adopted and deduced from the photometry by Castelli et al. (1985) ( $T_{\text{eff}}=14000$  K,  $\log g=4.0$ ), by Smith (1997) ( $T_{\text{eff}}=13990$  K,  $\log g=4.29$ ) and by van den Ancker et al. (1996) ( $T_{\text{eff}}=14000$  K,  $\log g=4.3$ ).

In this paper we examine the whole optical spectrum of HR 6000 from 3050 Å to 9460 Å with the aim to extend the analysis performed on IUE spectra by Castelli et al. (1985) to the visible region and to investigate the possible contamination of the stellar spectrum by a close T Tau companion. We re-examine the parameter determination, the line identification, and the abundances. We also searched for the presence of peculiarities like emission lines and isotopic anomalies which were recently discovered in HgMn stars. Owing to the availability of spectra taken at different epochs we also investigated possible spectral variabilities.

The comparison of the observed and computed spectra as described in this paper is available at the web-address given in the footnote<sup>1</sup>.

## 2. Observations and Radial Velocity

Spectra of HR 6000 were recorded at ESO on 2005 September 19 (JD 2453632.53881) and on 2006 March 25 (JD 2453819.90226). In both epochs we used the UVES DIC1 and DIC2 standard settings to cover the spectral range from 3050 Å to 9500 Å. The slit width was set to 0.3 for the red arm, corresponding to a resolving power of  $\lambda/\Delta\lambda \approx 110000$ . For the blue arm, we used a slit width of 0.4 to achieve a resolving power of  $\approx 80000$ . The spectra were reduced by the UVES pipeline Data Reduction Software (version 2.5; Ballester et al. 2000) and using standard IRAF routines. The signal-to-noise ratio (S/N) of the UVES spectra is very high, ranging from 150 in the near UV (3300 Å) to about 350 and 300 at 5000 Å and 7100 Å, respectively.

There are two gaps in the observed range at  $\lambda\lambda$  4523–4769 Å and 7536–7660 Å, which are caused by the physical gap between the two detector chips of the red CCD mosaic. In addition, the reduction in the spectral range 8076–8093 Å is affected by the presence of a bad column on the MIT CCD.

The heliocentric radial velocities derived from spectra observed on September 2005 and on March 2006 are  $2.6 \text{ km sec}^{-1}$  and  $1.4 \text{ km sec}^{-1}$ , respectively. These values are consistent with the mean heliocentric radial velocity  $+2.6\pm1.8 \text{ km s}^{-1}$  of the Lupus cloud (James et al., 2006), so that they strengthen the assumption of the membership of HR 6000 to the Lupus cloud. However, Catanzaro et al. (2004) measured  $0.67\pm0.38 \text{ km s}^{-1}$ , while Andersen & Jaschek (1984) determined  $-1.5\pm0.5 \text{ km s}^{-1}$ . In their study, the values obtained from different spectra range from  $+0.6\pm0.8 \text{ km s}^{-1}$  to  $-3.3\pm0.8 \text{ km s}^{-1}$ . Such differences in measured radial velocities by different authors indicate that small radial velocity variations in HR 6000 are possibly real.

The spectra were normalized to the continuum by using a modified version of the interactive code NORMA written by Bonifacio (1989). In the original procedure smoothed interpolated curves through selected continuum points are obtained by means of Hermite spline functions (subroutine INTEP by Hill

(1982)). We replaced the spline functions with a linear interpolation because in some other previous stellar analyses performed by one of us (F.C.) it has given a continuum more close to that we would have drawn by hand. However, we checked that in the case of HR 6000 the continua from the two different interpolations do not differ in a significant way. When needed, the continuum was adjusted over 6 Å intervals with the help of the synthetic spectrum. The continuous level just longward of the Balmer and Paschen discontinuities is highly uncertain because it is difficult to know where to put it. Also the continuum drawn over the broad hydrogen lines is uncertain owing to the not fully corrected distortions of the echelle spectra.

## 3. Stellar parameters

Table 1 summarizes the stellar parameters as determined from Strömgren, UBV, and near-infrared IJ photometry. The observed Strömgren and UBV indices were taken from the Hauck & Mermilliod (1998) Catalogue<sup>2</sup>, while  $(V-I_c)=-0.05$  was taken from the Hipparcos Catalogue and  $(V-J)=-0.032$  was obtained from the J magnitude given by Thé et al. (1996). The parameter determination is the result of an interpolation of the observed dereddened indices in the grid of synthetic indices based on the NEWODF model atmospheres (Castelli & Kurucz, 2004) computed for metallicity  $[M/H]=0.0$  and microturbulent velocity  $\xi=0.0 \text{ km s}^{-1}$ . The color grids are available at the web-address given in the footnote<sup>3</sup>. For the Strömgren and UBV photometry the adopted grids were uvbybeta and UBV, while for the  $(V-I_c)$  and  $(V-J)$  indices the adopted grid was UBVRJHKL.

Results from the Strömgren photometry are listed in the first panel of Table 1. For  $E(B-V)=0.06$ , the parameters from the  $c_0$  and  $\beta$  indices are  $T_{\text{eff}}=13827\pm200$  K,  $\log g=4.27\pm0.05$ . The reddening  $E(B-V)=0.06$  was determined by Bessell & Eggen (1972) and later confirmed by Thé & Tjin A Djie (1978). An uncertainty in  $E(B-V)$  of 0.02 mag corresponds to a difference in  $T_{\text{eff}}$  of only 15 K. The reddening  $E(B-V)=0.042$  listed in the first row of Table 1 corresponds to  $E(b-y)=0.031$  as yielded by the UBVLIST code of Moon (1985) on the basis of standard reddening relations. The conversion  $E(b-y)=0.74E(B-V)$  from Crawford & Mendwewala (1976) was applied here.

A microturbulent velocity of  $2.0 \text{ km s}^{-1}$  lowers the temperature by 100 K and does not affect the gravity, while an increase in metallicity to  $[M/H]=+0.5$  decreases  $T_{\text{eff}}$  by about 50 K; a decrease in metallicity to  $[M/H]=-1.0$  increases  $T_{\text{eff}}$  by about +300 K. The corresponding differences in  $\log g$  are of the order of 0.01 dex in all the cases. These values are only indicative of the metallicity effect in that the abundance pattern of HR 6000 is very different from any scaled solar one.


The parameters from the UBV photometry and  $\beta$  index are listed in the second panel of Table 1. For  $E(B-V)=0.06$ ,  $[M/H]=0.0$ , and  $\xi=0.0 \text{ km s}^{-1}$  they are  $T_{\text{eff}}=13973\pm300$  K,  $\log g=4.38\pm0.2$  (Fig. 1). They are close to those obtained from the Strömgren photometry. This parameter determination, described in Castelli et al. (1985), is very sensitive to small reddening changes. In fact, for  $E(B-V)=0.04$  or  $E(B-V)=0.08$  there is no intersection in the curves  $(U-B)_0$ ,  $(B-V)_0$  and  $\beta$ . For B-type stars, the  $(U-B)$ ,  $(B-V)$ , and  $\beta$  indices do not depend on the metallicity in a significant way (Castelli, 1999; Castelli & Kurucz, 2006).

The third panel of Table 1 shows that for a prefixed gravity  $\log g=4.0$ , the observed  $(V-I)_c$  and  $(V-J)$  indices are repro-

<sup>1</sup> <http://wwwuser.oat.ts.astro.it/castelli/hr6000/hr6000.html>

<sup>2</sup> <http://www.unige.ch/gcpd/gcpd.html>

<sup>3</sup> <http://wwwuser.oat.ts.astro.it/castelli/colors.html>



**Fig. 1.** Determination of the atmospheric parameters ( $T_{\text{eff}}$ ,  $\log g$ ) from  $(U-B)_0$ ,  $(B-V)_0$  and  $\beta$ . The circle represents the parameters  $T_{\text{eff}}=13873$  K,  $\log g=4.38$  obtained from this method for  $[M/H]=0.0$  and  $\xi=0.0$  km s $^{-1}$ . The observed indices were dereddened for  $E(B-V)=0.060$

duced by  $T_{\text{eff}}$  equal to 10889 K and 11517 K, respectively, when  $[M/H]=0.0$ ,  $\xi=0.0$  km s $^{-1}$ , and  $E(B-V)=0.06$ . We used the reddening relations  $E(V-I_c)=1.25E(B-V)$  from Dean et al. (1978) and  $E(V-J)=2.23E(B-V)$  from Mathis (1990). The temperature was derived by interpolating the observed dereddened  $(V-I_c)_0$  and  $(V-J)_0$  indices in the Bessell et al. (1998) color grid computed for  $[M/H]=0.0$  and  $\xi=0.0$  km s $^{-1}$ .

Catanzaro et al. (2004) were the first ones who derived parameters for HR 6000 from high resolution spectra, in particular from  $H_\gamma$  and  $H_\delta$  Balmer profiles observed on FEROS spectra. They fixed  $T_{\text{eff}}=12950\pm50$  K and  $\log g=4.05\pm0.01$  from ATLAS9 models computed for  $[M/H]=-0.5$  and zero helium abundance. This particular grid of models yields the same parameters from both Balmer profiles, but we note that the parameters do not change if the grid computed with  $[M/H]=0.0$  and solar He abundance is used, provided that the different parameters from  $H_\gamma$  ([12750 K, 3.95] and  $H_\delta$  ([13150 K, 4.13]) are averaged. As discussed in Sect. 4, parameters very close to those of Catanzaro et al. (2004) were derived by us from Balmer profiles and Fe I, Fe II equivalent widths measured on UVES spectra and from model atmospheres computed for the individual abundances of HR 6000.

The last panel of Table 1 shows that the cooler model, as adopted by Catanzaro et al. (2004) (model T12950g405k2M05nohe), does not reproduce the observed Strömgren indices. In fact, the difference between the observed and computed  $(b-y)_0$  indices amounts to 0.020 mag, or even more, depending on the adopted reddening, while the difference between the  $c_0$  indices is 0.122 mag. These discrepancies do not change in a remarkable way for the ATLAS12 model (model T12850g41k0at12) adopted by us to predict the observed spectrum (see Sect. 4). On the other hand, the hotter model [13900, 4.32] fails to reproduce all the Balmer profiles. For solar abundances it provides good agreement between observations and computations for  $H_\alpha$ , but the agreement decreases from  $H_\beta$

to  $H_\delta$  in the sense that the computed wings become too broad as compared with the observed ones; for individual abundances the computed inner wings of all the profiles are narrower than the observed ones, in particular for  $H_\alpha$  whose profile seems also to be a little bit asymmetric with a more extended red wing. All the comparisons between the observed and computed Balmer profiles are available at the website of footnote 1. Fig. 2 shows the case of  $H_\beta$ .

In conclusion, there is a discrepancy of about 1000 K between parameters derived from UBV and Strömgren photometry and parameters derived from the line spectrum. Furthermore, the difference in  $T_{\text{eff}}$  from the near-IR photometry is so large that spectrophotometric observations are needed in order to confirm these results. Unfortunately, the only spectrophotometric data available for HR 6000 are the IUE spectra. For  $E(B-V)=0.06$ , the best fit of the UV energy distribution from the IUE images SWP14849 and LWR11431 to the grid of ATLAS9 energy distributions computed for  $[M/H]=0.0$  and  $\xi=0.0$  km s $^{-1}$  is achieved with  $T_{\text{eff}}=13300$  K,  $\log g=4.1$ . More details about these IUE images can be found in Thé et al. (1996).

#### 4. The ATLAS12 model and abundance determination

The opacity distribution function ATLAS9 model with parameters derived from the UBV and Strömgren photometry ( $T_{\text{eff}}=13900$  K,  $\log g=4.32$ ,  $[M/H]=0.0$ ,  $\xi=0.0$  km s $^{-1}$ ) was used to derive preliminary abundances. Abundances for iron and phosphorus were obtained from equivalent widths, while those for all the other elements were obtained with the synthetic spectrum method. Mostly the lines listed in Castelli & Hubrig (2004b) (electronic Appendix) were examined. Synthetic spectra were computed with the SYNTH code (Kurucz, 1993). The adopted line lists are available on the web-address given in the footnote<sup>4</sup>. The synthetic spectra were broadened for a rotational velocity  $v \sin i=1.0$  km s $^{-1}$  and an instrumental resolving power of 80000 for  $\lambda < 4520$  Å and of 110000 for  $\lambda > 4765$  Å (Sect. 2). The adopted  $v \sin i$  was that reproducing the observed profiles at best, in particular Mg II 4481 Å. It is in close agreement with the value of 0.0 km s $^{-1}$  derived by Catanzaro et al. (2004).


With this model, the average abundance derived from the equivalent widths of 11 Fe I selected lines was equal, within the errors limits, to that obtained from the equivalent widths of 34 selected Fe II lines, i.e.  $\log(N(\text{Fe I})/N_{\text{tot}})=-3.65\pm0.06$  and  $\log(N(\text{Fe II})/N_{\text{tot}})=-3.69\pm0.15$ . At the same time, for  $\xi=0.0$  km s $^{-1}$  there was no trend of the abundances versus the equivalent widths of the Fe II lines. The table with the measured equivalent widths and corresponding abundances is available at the website of footnote 1.

The abundances based on the ATLAS9 model were used as input data for an opacity sampling ATLAS12 model (Kurucz 1997; Castelli & Kurucz, 1994) computed for the same parameters of the ATLAS9 model. Due to the extremely peculiar abundances of HR 6000, the  $T$ - $\tau_{\text{Ross}}$  structure of the ATLAS12 model is rather different from that of the ATLAS9 model, in particular in layers upper than  $\log(\tau_{\text{Ross}})\leq -0.5$ . The comparison of the  $T$ - $\tau_{\text{Ross}}$  relations from ATLAS9 and ATLAS12 is similar to that shown in Castelli & Hubrig (2004b). As a consequence, in spite of the achieved better agreement between observed and computed Balmer profiles, the Fe I-Fe II ionization equilibrium worsens. The average abundances for Fe I and Fe II became  $-3.42\pm0.06$  dex and  $-3.61\pm0.15$  dex, respectively.

<sup>4</sup> <http://wwwuser.oat.ts.astro.it/castelli/linelists.html>

**Table 1.** Model parameters for HR 6000 from photometry by assuming  $[M/H]=0.0$  and  $\xi=0.0 \text{ km s}^{-1}$ .

Observed indices				Dereddened indices				Parameters		
(b-y)	$m_1$	$c_1$	$\beta$	E(b-y)	(b-y) <sub>0</sub>	$m_0$	$c_0$	$T_{\text{eff}}$ (K)	$\log g$	E(B-V)
-0.030	0.116	0.512	2.750	0.031	-0.061	0.126	0.506	13799	4.27	0.042
$\pm 0.003$	$\pm 0.003$	$\pm 0.003$	$\pm 0.013$	0.044	-0.074	0.130	0.503	$\pm 150$	$\pm 0.05$	
								13827	4.27	0.060
								$\pm 200$	$\pm 0.05$	
V	U-B	B-V		E(B-V)	$V_0$	(U-B) <sub>0</sub>	(B-V) <sub>0</sub>	$T_{\text{eff}}$ (K)	$\log g$	
6.647	-0.434	-0.073		0.06	6.461	-0.477	-0.133	13973	4.38	
$\pm 0.013$	$\pm 0.004$	$\pm 0.004$						$\pm 300$	$\pm 0.2$	
(V-I) <sub>c</sub>		(V-J)		E(V-I) <sub>c</sub>	E(V-J)	(V-I) <sub>0</sub>	(V-J) <sub>0</sub>	$T_{\text{eff}}$ (V-J)	$T_{\text{eff}}$ (V-I) <sub>c</sub>	$\log g$
-0.05		-0.032		0.075	0.134	-0.125	-0.166	11517	10889	4.0 (prefixed)
$\pm 0.00$										
model				(b-y) <sub>calc</sub>		$c_{\text{calc}}$				
T12950G405k2M05nohe				-0.041		0.629				
T12850g41k0at12				-0.049		0.596				

**Fig. 3.** Comparison of IUE data (dotted line) with the flux computed from the ATLAS12 model with individual stellar abundances (full line). Model parameters are  $T_{\text{eff}}=12850 \text{ K}$ ,  $\log g=4.1$ , and  $\xi=0.0 \text{ km s}^{-1}$ . The IUE flux was dereddened for  $E(B-V)=0.06$ .

A search for an appropriate ATLAS12 model suitable to reproduce both the Balmer profiles and the Fe I-Fe II ionization equilibrium has finally yielded, after several trials and errors, the parameters  $T_{\text{eff}}=12850 \text{ K}$ ,  $\log g=4.1$ , in good agreement with the values adopted by Catanzaro et al. (2004). The Fe I and Fe II abundances were  $-3.89 \pm 0.06 \text{ dex}$  and  $-3.88 \pm 0.16 \text{ dex}$ , respectively. Fig. 3 shows that this model also reproduces, for  $E(B-V)=0.06$ , the slope of the UV energy distribution given by the IUE images SWP14849 and LWR11431. However, a close inspection of Fig. 3 reveals the presence of some strong absorptions in the short-wavelength region which are neither predicted by the models nor detected in the examined IUE high-resolution spectra (Sect. 5.1). As already discussed in the previous section, the final ATLAS12 model fails in reproducing the observed Strömgren indices. The conclusion is that we were not able to fix a model well suited to reproduce the atmosphere of HR 6000 as deduced from the available observations. Table 2 summarizes the parameter values from the different determinations based on the ATLAS9 (AT9) and ATLAS12 (AT12) models.

**Table 2.** Summary of the stellar parameters for HR 6000 from different determinations.

Method	$T_{\text{eff}}$	$\log g$	Models
UBV $\beta$	$13973 \pm 300$	$4.4 \pm 0.2$	AT9
$c_0\beta$	$13813 \pm 200$	$4.3 \pm 0.05$	AT9
IUE flux	$13300 \pm 100$	$4.1 \pm 0.1$	AT9
Balmer profiles	$12850 \pm 50$	$4.1 \pm 0.01$	AT12
Fe I/Fe II	12850	4.1	AT12
VI <sub>c</sub> J	$11203 \pm 300$	4.0 (prefixed)	AT9

## 5. The resulting abundances


The abundances used for computing the whole synthetic spectrum of HR 6000 from  $3050 \text{ Å}$  to  $9460 \text{ Å}$  are listed in column 2 of Table 3. The synthetic spectrum overimposed on the observed spectrum is available at the website of footnote 1. The abundances are based on the ATLAS12 model with parameters  $T_{\text{eff}}=12850 \text{ K}$ ,  $\log g=4.1$ ,  $\xi=0.0 \text{ km s}^{-1}$ , and  $v \sin i=1.0 \text{ km s}^{-1}$ . They are shortly discussed below.

The main results are the large underabundance of Si ( $[-2.9]$ ) followed by that of C ( $[-1.8]$ ). Other underabundant elements are He ( $\leq [-0.85]$ ), N ( $\leq [-1.7]$ ), O ( $[-0.5]$ ), Ne ( $[-0.9]$ ), Mg ( $[-1.2]$ ), Al ( $[-1.7]$ ), S ( $[-1.5]$ ), Cl ( $< [-0.7]$ ), Ca ( $[-0.2]$ ), Sc ( $[-0.6]$ ), V ( $[-1.1]$ ), Co ( $\leq [-0.8]$ ), Ni ( $[-0.4]$ ), and Sr ( $[-1.0]$ ).

On the other hand, the most overabundant element is Xe ( $[+4.5]$ ), followed by Hg ( $[+2.7]$ ), and P ( $[+2.3]$ ). Other overabundant elements are Be ( $[+0.7]$ ), Ti ( $[+0.25]$ ), Mn ( $[+0.95]$ ), Fe ( $[+0.7]$ ), Y ( $[+1.0]$ ), and possibly Ba ( $[+1.0]$ ).

Below we discuss the individual abundances:

**Helium:** From the analysis of the He I lines  $4713 \text{ Å}$  and  $5875 \text{ Å}$ , Catanzaro et al. (2004) derive an underabundance of  $[-1.45]$ , i.e.  $N(\text{He})/N_{\text{tot}}=0.0032$ . We extended their analysis to all the He I lines previously analyzed in the HgMn star HD 175640 by Castelli & Hubrig (2004b), except for He I  $4713 \text{ Å}$  which is located in a gap of the observed spectral regions. While all the He I lines were fairly well predicted in HD 175640 by the same abundance, this is not a case in HR 6000. In fact, the abundance  $N(\text{He})/N_{\text{tot}}$  from the different lines ranges from 0.0125 to 0.002, i.e. from  $[-0.80]$  to  $[-1.60]$ . The higher value reproduces the whole profile of He I  $3867.4 \text{ Å}$



**Fig. 2.** The observed  $H_\beta$  (thick line) is compared with profiles computed with different models (thin lines). In the upper and middle panels the ATLAS9 model and the ATLAS12 model with parameters given by the photometry ( $T_{\text{eff}}=13900\text{ K}$ ,  $\log g=4.32$ ,  $\xi=0.0\text{ km s}^{-1}$ ) were used. The lower panel shows the profile computed with the final adopted ATLAS12 model with parameters  $T_{\text{eff}}=12850\text{ K}$ ,  $\log g=4.1$ , and  $\xi=0.0\text{ km s}^{-1}$ .


and only the wings of He I lines at  $4026.1\text{ \AA}$  and  $4387.9\text{ \AA}$ , whose computed cores appear stronger than the observed ones (Fig. 4). The same discrepancy occurs also for the lines at  $\lambda\lambda$  4471.4, 4921.9, 5015.7  $\text{\AA}$ , whose wings are fitted by abundances equal to 0.005 and 0.006, while the cores would require a lower value. The two weak lines at  $\lambda\lambda$  5875.6 and 6678  $\text{\AA}$  are rather well predicted for  $N(\text{He})/N_{\text{tot}}$  equal to 0.002 and 0.003, respectively. Finally, the line at  $7065.2\text{ \AA}$  is partially blended both with telluric lines and with Fe II 7065.163  $\text{\AA}$ . It can not be reproduced whichever is the abundance. The abundance from the remaining He I lines, which are very weak, depends on the assumed position for the continuum level. The comparison between observed and computed He I profiles is available at the website of footnote 1.

Profiles of He I like those observed in HR 6000 were observed also in other HgMn stars as for instance HR 7361, HR 7664 (Dworetzky, 2004) or Feige 86 (Bonifacio et al., 1995). This kind of anomalous profiles was interpreted as evidence of helium stratification (Dworetzky, 2004; Bohlender, 2005).

No contribution of  $^3\text{He}$  to  $\lambda$  6678  $\text{\AA}$  was observed.

**Beryllium:** The abundance was derived from the lines at 3130.42 and 3131.06  $\text{\AA}$ , although the second one is heavy blended.

**CNO:** While Catanzaro et al. (2004) did not observe C II lines, we possibly identified very weak C II lines at  $\lambda\lambda$  3918.968,



**Fig. 4.** Comparison of observed (thick line) and computed (thin line) He I 4026  $\text{\AA}$  profiles. The abundance  $\log(N(\text{He})/N_{\text{tot}})=-1.90$  which fits the wings yields a too strong computed core.

**Table 3.** Final abundances  $\log(N_{\text{elem}}/N_{\text{tot}})$  from the ATLAS12 model with parameters  $T_{\text{eff}}=12850\text{ K}$ ,  $\log g=4.1$ ,  $\xi=0.0\text{ km s}^{-1}$ . Solar abundances are from Grevesse & Sauval (1998).

Elem	UVES	IUE(old)	IUE(new)	Sun
He	-1.90			-1.05
Be	-9.90	$\leq -9.5$	-9.90	-10.64
B		-10.1	-10.29	-9.49
C	-5.32:	$\geq -5.4$	-5.82	-3.52
N	$\leq -5.82$	$\geq -6.5$	-6.50	-4.12
O	-3.71	-3.3	-3.70	-3.21
Ne	-4.86			-3.96
Na	-5.71			-5.71
Mg	-5.66	-5.6	-5.66	-4.46
Al	-7.30	-7.7	-7.30	-5.57
Si	-7.40	$< -5.7$	-7.40	-4.49
P	-4.30	-4.4--5.0	-5.00	-6.59
S	-6.20	-6.0--5.5	-6.20	-4.71
Cl	$\leq -7.24$	-4.3	-7.24	-6.54
Ca	-5.88	-5.2	-5.88	-5.68
Sc	-9.50	-11.0	-9.50	-8.87
Ti	-6.77	$> -6.5$	-6.77	-7.02
V	-9.14	-6.7--8.0	-9.14	-8.04
Cr	-6.37	-6.2	-6.37	-6.37
Mn	-5.60	-5.3	-5.60	-6.65
Fe	-3.85	-3.8	-3.85	-4.54
Co	$< -7.92$	$\geq -9.3$	-7.92	-7.12
Ni	-6.19	-7.0	-6.19	-5.79
Cu	$\leq -7.83$	$\geq -9.3$	-7.83	-7.83
Zn		-8.3	-7.44	-7.44
Ga	$\leq -9.16$	-8.4	-9.16	-9.16
Sr	-10.07			-9.07
Y	-8.80			-9.80
Xe	-5.40			-9.87
Ba	-8.91?			-9.91
Hg	-8.20	-	-8.20	-10.91

4267.01, and 4267.26 Å. No nitrogen lines were observed. We deduced a carbon underabundance of  $[-1.80]$  and an upper limit for nitrogen of  $[-3.40]$ . Oxygen is underabundant by  $[-0.50]$ . We examined all the O I lines listed in Castelli & Hubrig (2004b). The O I infrared lines at  $\lambda\lambda$  7774, 8446, and 9260 Å are inadequately computed under the LTE assumption.

*Neon:* The abundance was fixed from Ne I 7032.413 Å.

*Sodium:* There is a remarkable inconsistency between the abundance of  $-3.31$  dex from the two lines Na I at 3302.368 Å and 3302.978 Å and the almost solar abundance of  $-5.57$  dex of the Na I lines at  $\lambda\lambda$  5688.2, 8183.2 and 8194.8 Å. As already described by Catanzaro et al. (2004) the two stellar Na I D lines at 5889.951 Å and 5895.924 Å are completely hidden by the interstellar lines. The most likely explanation is that the lines at 3302 Å are also of interstellar origin. They have also been observed in the spectrum of HD 175640 as weak red-shifted absorptions with the same wavelength displacement of  $+0.08$  Å as the interstellar D-lines.

*Magnesium:* The estimated underabundance is  $[-1.20]$ . Magnesium abundance was mostly derived from Mg II 4481 Å. The Mg II doublet at 6545.942, 6545.994 Å is much broader than predicted and is also red-shifted. The most plausible explanation is the lack of Stark broadening data in the computations.

*Aluminium:* The estimated underabundance is  $[-1.73]$  from the Al II lines at  $\lambda\lambda$  7042.06 and 7056.60 Å. No other lines, as Al II at  $\lambda\lambda$  3944.009 and 3961.523 Å, were observed.

*Silicon:* The estimated underabundance is  $[-2.90]$ . Only weak Si II lines at  $\lambda\lambda$  3853.665, 3856.018, and 3862.595 Å were observed.

*Phosphorus:* The overabundance of phosphorus can be inferred from most of the P II and P III lines, but the real value is very difficult to be deduced because the different lines give rather different abundances. We measured the equivalent widths for a sample of 22 P II and 4 P III lines. The average abundance from P II is  $\log(N(\text{P})/N_{\text{tot}}) = -4.36 \pm 0.28$  for  $\xi = 0.0\text{ km s}^{-1}$ , but the trend of the abundances with the equivalent widths is very large. It disappears for  $\xi = 6.0\text{ km s}^{-1}$ . The corresponding abundance of  $-4.70 \pm 0.12$  produces, for  $\xi = 6.0\text{ km s}^{-1}$ , extremely broad and shallow profiles completely different from the observe ones.

The abundance for  $\xi = 0.0\text{ km s}^{-1}$  inferred from the P III lines is  $\log(N(\text{P})/N_{\text{tot}}) = -4.57 \pm 0.27$ . Therefore, the P II abundance for  $\xi = 0.0\text{ km s}^{-1}$  is 0.2 dex larger than that obtained from P III, but lies within the large mean square error of the averages. The large uncertainty in the phosphorous abundance could be related to NLTE effects which have been neglected here. Another possibility could be the presence of phosphorus vertical abundance stratification.

The adopted P II and P III lines, together with the atomic data and the measured equivalent widths, are listed on the website of footnote 1.

*Sulfur:* The sulfur underabundance of  $[-1.5]$  was derived from the same lines used by Castelli & Hubrig (2004b).

*Chlorine:* The absence of observed chlorine lines, like Cl II 4794.54 Å, implies an abundance less than  $-7.24$  dex.

*Potassium:* Potassium can be predicted by adopting solar abundances. However, the slight blue-shift of  $-0.05$  Å of the lines K I  $\lambda\lambda$  7664.911, 7698.974 Å, the only ones observed in the spectrum, may indicate an interstellar origin for them.

*Calcium:* All the Ca II lines listed in Castelli & Hubrig (2004b) can be reproduced by an underabundance of  $[-0.2]$ , except for the lines of mult. 13 (8201.720 Å and 8248.796 Å) which are predicted too strong. The blue wing of the K line at 3933.664 Å presents an additional broad weak absorption. It is further discussed in Sect. 6.3. The lines of the Ca II infrared triplet are red-shifted by 0.14 Å. They are discussed in more detail in Sect. 6.1.

The Ca I line at 4226.728 Å is very weak. It is reproduced by solar abundance. A possible interstellar contribution can not be excluded.

*Scandium:* The underabundance of  $[-0.6]$  was derived from the Sc II lines listed in Castelli & Hubrig (2004b).

*Titanium:* The overabundance of  $[+0.25]$  was derived from the Ti II lines listed in Castelli & Hubrig (2004b).

*Vanadium:* The underabundance of  $[-1.1]$  was deduced from the lines of V II mult. 1 at 3100 Å.

*Chromium:* Most of the observed Cr II lines can be predicted by solar abundance.

*Manganese:* The abundance of  $-5.0$  dex from the lines of Mn II mult. 3 at 3440–3498 Å is larger by 0.6 dex than that obtained from most of the Mn II lines lying longward of the Balmer discontinuity. This result indicates the possible occurrence of vertical stratification for the abundance of this ion. The abundance of  $-5.6$  dex listed in Table 3 is the one obtained from the region longward of the Balmer discontinuity. Its value is the same as that presented by Catanzaro et al. (2004).

*Iron:* The abundance was derived from the equivalent widths of 11 Fe I and 34 Fe II lines. The lines together with the atomic data and the measured equivalent widths are available on the website of footnote 1. The overabundance of  $[+0.7]$  gives rise

in the spectrum to a very large number of Fe I, Fe II, and especially Fe II lines (see sub-section 6.4).

**Cobalt:** No Co II lines were observed in the spectrum, in particular the line at 3501.717 Å is absent. Therefore only an upper limit of  $[-0.8]$  was fixed.

**Nickel:** Most Ni II lines indicate an underabundance of  $[-0.4]$ .

**Copper:** No lines of Cu II were observed, in particular those at  $\lambda\lambda$  4909.734, 4917.892, and 4931.698 Å. Because for solar abundance there are no predicted lines we assumed solar abundances as upper limit for this element.

**Gallium:** No gallium lines were observed. If a feature at  $\lambda$  5416.32 Å is identified as Ga II it would be reproduced by  $\log(N(\text{Ga})/N_{\text{tot}}) = -6.0$ . This abundance gives rise to other strong computed Ga II lines with no observed counterparts.

**Strontium:** No Sr II lines were observed. Only a very weak absorption is detectable at the position of Sr II 4077.71 Å. It is reproduced by assuming an underabundance of  $[-1.0]$ .

**Yttrium:** The overabundance of  $[+1.0]$  was derived mostly from the Y II line at 4900.12 Å.

**Zirconium:** No Zr lines were observed in the spectrum. They are not predicted by assuming solar Zr abundance.

**Xenon:** Xenon is the most overabundant element  $[+4.5]$ . The abundance was derived from the Xe II lines listed in Castelli & Hubrig (2004b) having  $\log gf$  available from the NIST database. They were taken from Wiese & Martin (1980) (WM80). Furthermore, we added in the line list some lines with  $\log gf$ 's from Ryabchikova & Smirnov (1989). In addition, only for identification purposes, all the lines with no  $\log gf$  in the NIST database but with intensity  $\geq 100$  were included in the line lists. A guessed  $\log gf$  was assigned to them.

We noticed that the position of several Xe II lines is close to that of unidentified stellar lines with the wavelength blue-shifted by 0.05 to 0.1 Å from the Xe II laboratory wavelength. Examples are the lines at  $\lambda\lambda$  4180.10, 4208.48, 4209.47, 4238.25, 4245.38, 4330.52, 4393.20, 4448.13, 4462.19, 5667.56 Å. We have observed the same occurrence also in the two Xe rich HgMn stars Feige 86 and 46 Aql (Castelli & Hubrig, in preparation). If the Xe II identification is correct, the wavelength shift could be due to some isotopic anomaly. In fact, no isotopic composition was considered in our computations owing to the lack of isotopic wavelengths for Xe II. There are data only for  $\lambda$  6051.150 Å (Alvarez et al., 1979). For this line the isotopic wavelengths range from 6051.148 Å to 6051.152 Å. Because the stellar line is blue-shifted by 0.03 Å from the laboratory wavelength 6051.150 Å (Hansen & Persson, 1987) some isotopic anomaly can not be invoked to explain the observed shift.

**Barium:** We were not able to draw any conclusive results for barium. There is a weak absorption at 4933.93 Å, but if we identify it as Ba II 4934.076 Å, this barium line would be blue-shifted by 0.15 Å. The wavelength in the spectrum would correspond to the Ritz wavelength 4933.970 Å rather than to the observed wavelength 4934.077 Å, as they are listed in the NIST database. Any isotopic anomaly has to be excluded because the shortest isotopic wavelength is that of the hyperfine component of  $^{137}\text{Ba}$  at 4934.054 Å (McWilliam, 1998).

**Mercury:** Only Hg II at 3984 Å is observable. It can be predicted by assuming an overabundance of  $[+2.7]$  and a non-terrestrial isotopic composition. The Hg isotopic anomaly is discussed in Sect. 6.1.

## 5.1. Comparison with the abundances from the IUE spectra

Column 3 of Table 3 lists the abundances that were derived from IUE spectra by Castelli et al. (1985). The analyzed images were SWP04588 and LWR03979. We revised these abundances by using the same ATLAS12 model adopted for the analysis of the UVES spectra and the same images that we downloaded from the MAST archive<sup>5</sup>. Also updated line lists, as compared with those used by Castelli et al. (1985), were used. Column 4 of Table 3 lists the revised abundances from the ultraviolet.

The IUE spectra are very noisy and the resolving power is too low ( $R \sim 12000$ -13000) to allow one to perform an accurate analysis. Almost all very numerous lines are blended either with known or with unknown components. In addition, the never adequately corrected distortions of the echelle orders make the drawing of the continuum a very problematic task. The large uncertainty in the continuum position affects both the shape and the strength of the spectral lines. For all these reasons the abundances that we derived from the IUE spectra have always been considered as estimates rather than definitive values.

Table 3 shows that the difference IUE(new)–IUE(old) between revised and Castelli et al. (1985) abundances is larger than 0.4 dex for C, O, Cl, Ca, Sc V, Ni, Zn, and Ga. The abundances from UVES spectra reproduce the IUE spectrum (within all the above outlined limits) for all the elements except for carbon and phosphorous, for which the differences UVES–IUE(new) are +0.5 dex and +0.7 dex respectively. While the analysis of the carbon abundance from the visible spectral region is based only on very weak blended lines and therefore is rather uncertain, the phosphorous discrepancy confirms the difficulty in fixing the phosphorous abundance from the different lines.

The synthetic spectrum overimposed on the IUE spectra is available at the website of footnote 1. It was computed with the abundances listed in column 4 of Table 3.

## 6. Line spectrum peculiarities

In the following sub-sections we discuss isotopic anomalies observed for Hg and Ca and the emissions observed for a few lines of Mn II, Cr II, and Fe II. We also discuss the huge quantity of lines which could not be identified.

### 6.1. Isotopic anomalies

Isotopic anomalies in HR 6000 were observed for the Ca II infrared triplet and for Hg II at 3984 Å. No  $^3\text{He}/^4\text{He}$  isotopic anomaly was detected in the UVES spectra.

**Ca II infrared triplet:** The lines of the infrared triplet at  $\lambda\lambda$  8498.023, 8542.091, and 8662.141 Å are red-shifted by 0.14 Å and appear broader and stronger than the predicted ones. The wavelength red-shift of the infrared triplet observed in several CP stars (Cowley et al., 2007) was interpreted by Castelli & Hubrig (2004a) as due to a Ca isotopic composition different from the terrestrial one.


**Hg II 3984:** Hg II at 3984 Å, although weak, was detected in the UVES spectra. The isotopic composition that best reproduces the observed line for  $\log(N(\text{Hg})/N_{\text{tot}}) = -8.20$  is that listed in column 3 of Table 4. For comparison, the terrestrial isotopic composition (Anders & Grevesse 1989) is given in column 5 of Table 4. In HR 6000 the isotope 204 is the most abundant one. The profiles computed with the stellar and terrestrial isotopic

<sup>5</sup> <http://archive.stsci.edu/iue/>



**Table 4.** Mercury isotopic (iso) and hyperfine (hfs) composition in % for Hg II at 3984 Å.

Isotope	$\lambda$	stellar %	log $gf$ iso	terr. %	log $gf$ iso	hfs. %
196	3983.769	0.15	-2.81	0.15	-2.81	
198	3983.838	0.77	-2.11	9.97	-1.00	
199a	3983.838	20.00	-0.70	16.87	-0.77	7.14
199b	3983.849	20.00	-0.70	16.87	-0.77	9.71
200	3983.909	6.87	-1.16	23.10	-0.64	
201a	3983.930	5.00	-1.30	13.18	-1.32	4.80
201b	3983.941	5.00	-1.30	13.18	-1.08	8.30
202	3983.990	10.00	-1.00	29.86	-0.52	
204	3984.071	57.20	-0.24	6.87	-1.16	

**Fig. 5.** The observed profile (thick line) of Hg II 3984 Å is compared in the lower panel with the profile computed with a terrestrial isotopic composition (thin line) and in the upper panel with the profile computed with the isotopic mixture listed in column 3 of table 4 (thin line).

compositions are compared with the observed profile in Fig. 5. We note that the Hg II line is blended with Fe I 3983.956 Å.

Hg II at 1942.3 Å can be observed in the IUE spectra, but the isotopic shifts are too small to enable us to confirm the isotopic composition deduced from the line at 3984 Å.

## 6.2. Emission lines

Emission lines were observed for Mn II, Cr II, and Fe II. We assumed that the studied lines are true emissions only if they were

observed in both UVES spectra taken at two different epochs. Emission lines are listed in Table 5.

*Emission lines of Mn II:* Some lines of Mn II multiplet 13 have been observed either in emission ( $\lambda\lambda$  6125.863, 6126.218, 6126.510, 6131.923 Å) or much weaker than predicted ( $\lambda\lambda$  6122.434, 6122.810, 6128.734 Å). Some others lines ( $\lambda\lambda$  6129.033, 6130.796, 6131.016 Å) are not observed in the spectrum taken on September 2005, while they appear as weak emissions in the spectrum of March 2006. This fact may suggests the presence of some variability in the emissions. Finally, the two very weak Mn II lines of mult. 13 at 6123.160 Å and 6129.254 Å are in absorption and agree rather well with the predicted lines.

Other weak Mn II emissions occur at 6446.337 Å (Mn II, mult. 19) which is blended with Fe II 6446.410 Å, and at 7219.968, 8495.229, 8565.819, 8769.175 Å. An emission could also affect the line at 8695.208 Å, which is observed much weaker than predicted, and the line at 8819.584 Å which, although predicted, is not observed. We note that there is a rather good agreement for this last line between the observed and computed profiles in the HgMn star HD 175640 (Castelli & Hubrig, 2004b), even considering that no hyperfine structure is available for computations.

*Emission lines of Cr II:* A few Cr II lines arising from high excitation levels and having large log  $gf$ 's appear clearly in emission. These lines were also found in emission in HD 175640 (Castelli & Hubrig, 2004b).

*Emission lines of Fe II:* There are no clear emissions for Fe II. The most probable ones are listed in Table 5. However, there is a set of observed lines much weaker than predicted. This disagreement could be explained either due to incorrect log  $gf$ 's or to emissions filling the absorptions. They occur at  $\lambda\lambda$  6145.611, 6214.948, 6328.498, 6340.841, 6349.601, 6419.626, 6436.598, 6451.094, 6479.607, 6480.807, 6518.774, 6524.726, 6576.153, 6576.170 Å.

The phenomenon of weak emission lines at optical wavelengths for main-sequence B-type stars has been noted from observations of relatively few stars. Presently, explanations of this phenomenon have been put forward in the context of the non-LTE line formation (Sigut, 2001) and possible fluorescence mechanisms (Wahlgren & Hubrig, 2000).

## 6.3. Unidentified lines

There is a huge number of unidentified lines in the spectrum of HR 6000. Their list is available at the web-address given in footnote 1. For instance, most impressive regions are  $\lambda\lambda$  4404-4411 Å and 5100-5300 Å, which are overcrowded with unidentified absorptions. We estimate that most of them are due to Fe II. In fact, a comparison with the spectrum of HD 175640 (Castelli & Hubrig, 2004b) has shown unidentified lines at the same wavelengths also in this star, although less intense than in HR 6000, in accordance with the different iron abundances (Fig. 6). Therefore the unidentified lines are not a special peculiar characteristic of HR 6000.

In the Kurucz line list for Fe II (Kurucz, 2005)<sup>6</sup> there are numerous transitions between observed levels with wavelengths close to those of the unidentified lines, but with so low log  $gf$ 's that none corresponding absorption is predicted. The comparison of log  $gf$ 's from Kurucz with the log  $gf$ 's from Raassen & Uylings (1998) has shown agreement in the two determina-

<sup>6</sup> <http://kurucz.harvard.edu/atoms/2601/gf2601.pos>



**Table 5.** Emission lines observed in HR 6000.

$\lambda(\text{\AA})$	Element	Mult.	$\log gf$	$\chi_{low}$	Notes
5518.08	?				
6112.906	Fe II?	–	–3.753	90300.625	not predicted.
6122.434	Mn II	13	0.950	82136.400	absorption filled by emission?
6122.810	Mn II	13	0.084	82136.400	absorption filled by emission?
6125.863	Mn II	13	0.783	82144.480	
6126.218	Mn II	13	0.230	82144.480	
6126.510	Mn II	13	–0.791	82144.480	
6128.734	Mn II	13	0.588	82151.160	absorption filled by emission?
6129.033	Mn II	13	0.208	82151.160	variable?
6130.796	Mn II	13	0.354	82155.840	variable?
6131.016	Mn II	13	0.053	82155.840	blend with Fe II in absorption
6131.923	Mn II	13	0.053	82158.170	
6158.621	Cr II	–	0.718	89174.080	
6182.340	Cr II	–	0.452	89336.890	
6446.337	Mn II	19	–0.663	98423.300	on the blue wing of Fe II 6446.410 Å
6526.302	Cr II	–	0.173	89885.080	
6585.241	Cr II	–	0.829	90850.960	red-shifted
7174.368	Mn II	–	0.397	85960.460	
7175.815	Mn II	–	–0.023	85960.460	
7219.968	Mn II	–	0.768	86057.440	
7222.639	Mn II	–	0.252	86057.440	
8495.229	Mn II	–	0.198	85960.460	
8565.819	Mn II	–	0.278	86057.440	
8695.208	Mn II	–	0.577	74560.010	absorption filled by emission?
8740.349	Fe II	–	0.002	99068.450	
8758.410	?				
8769.175	Mn II	–	0.491	74560.010	
8819.584	Mn II	13	0.347	74560.010	absorption filled by emission?
8839.005	Fe II	–			J78, unclassified <sup>a</sup>
9020.127	Fe II	–			J78, unclassified <sup>a</sup>
9085.11	?				
9208.4	?				

<sup>a</sup>J78=Johansson(1978)

tions. Therefore, either both determinations are off by several orders of magnitudes or other transitions at the same wavelength occur. This possibility seems partly justified when the Kurucz line list including lines from both observed and predicted levels is used<sup>7</sup>. In fact, numerous Fe II lines due to transitions between predicted levels have  $\log gf$ 's large enough to produce observable absorptions. Unfortunately, the uncertainty in their wavelength may be as large as 1 Å and also the corresponding  $\log gf$ 's may be affected by rather large errors. Table 6 is an example of tentative line identification for the unidentified lines in the interval 5130–5136 Å. As it is shown in Fig. 6 and in the lower panel of Fig. 7, out of displayed 17 spectral lines only three lines can be identified as arising from observed levels. They are Fe II 5131.210 Å, Fe II 5132.669 Å, and Fe I 5133.688 Å. The middle panel of Fig. 7 shows that more lines would be identified if several  $\log gf$ 's listed in col. 5 of Table 6 are replaced by those given in col. 7. They were derived by fitting the computed lines to the observed lines for the stellar iron abundance of –3.85 dex. Upper panel of Fig. 7 shows that the number of identified lines increases if the Fe II lines from predicted levels listed in cols. 8 and 9 of Table 6 are added. The intensity of the lines from predicted levels is usually lower than that of the observed lines.

There is a remarkable coincidence between several Fe II lines measured by Johansson (1978) (Table 6, col. 3) and the unidentified lines in our spectra. However, they are not predicted

by the synthetic spectrum either because they are missing in the line lists or owing to their too low assigned  $\log gf$ .

## 7. The T Tauri companion

Van den Ancker et al. (1996) suggested the presence of a T Tauri companion for HR 6000 in order to explain the strong X-ray emission observed by Zinnecker & Preibisch (1994) and the infrared excess that they inferred from the comparison of the observed energy distribution with that computed for the best fitting model having parameters  $T_{\text{eff}}=14000\text{ K}$ ,  $\log g=4.3$ . They assumed a temperature of 3500 K for the T Tauri central star, an accretion rate of  $2 \times 10^{-7} M_{\odot} \text{ yr}^{-1}$ , and a V magnitude of about 13.5, so that the companion would be about 6 mag fainter than HR 6000 and can not contribute significantly to the observed visual spectrum.

There are no clear spectroscopic traces of a second star in the spectrum of HR 6000, except for a broad weak absorption at  $\lambda\lambda 6707\text{--}6708\text{ \AA}$  which, although of different intensity in the two spectra observed on September 2005 and March 2006, exhibits a very similar shape. The synthetic spectrum of HR 6000 considered as a single star is flat in this region except for two weak Fe II lines at 6707.232 Å and 6707.539 Å which do not cover the whole observed absorption (Fig. 8, lower panel). We could argue that the feature is Li I 6707.7 which is a key element for the classification of T Tauri stars (Krautter et al., 1997). If such a star is present, it should be similar to a weak emission line T Tauri

<sup>7</sup> <http://kurucz.harvard.edu/atoms/gf2601/gf2601.lines0600>



**Table 6.** Unidentified lines in the region 5130-5136 Å.

$\lambda(\text{obs})$ (Å)	Identif.	$\lambda$ J78 <sup>a</sup>	$\lambda$ K03 <sup>b</sup>	$\log gf$ K03 <sup>b</sup>	$\log gf$ RU <sup>c</sup>	$\log gf$ Modified	$\lambda$ predicted(K03) <sup>b</sup>	$\log gf$
5130.0	Fe II	5130.011	5130.013	-5.590	-5.358	-1.790	5130.006	1.172
5130.18	Fe II	5130.177	missing				5130.196	-0.673
	Fe II						5130.198	-0.573
5130.3	Fe II		5130.291	-6.352		-1.252		
5130.45	Fe II						5130.427	-0.568
5130.60	Fe II	5130.605 uncl.	5130.618	-6.012	-5.562	-0.312	5130.542	-0.120
5131.4	Fe II						5131.478	-0.570
5131.5	Fe II						5131.650	-0.508
5131.7	Fe II		5131.706	-5.127		-0.127	5131.878	-0.727
	Fe II		5131.788	-5.023		-0.723		
5132.0	Fe II		5132.002	-4.648	-4.590	-1.648	5131.953	-0.222
5132.55	Fe II						5132.554	-0.190
5133.59	Fe II		5133.589	-3.961		-1.061	5133.475	0.220
5134.07	Fe II		5134.082	-5.805		-0.105		
5134.2	?							
5134.9	?							
5135.25	Fe II	5135.268	missing					
	Fe II		5135.187	-6.671		-2.671		
	Fe II		5135.297	-5.119	-4.394	-1.719		
5135.35	?							
5135.95	?							

<sup>a</sup>J78=Johansson (1978)<sup>b</sup>K03=Kurucz (2003)<sup>c</sup>RU=Raassen & Uylings (1998)

of  $0.1 \pm 0.2$  dex,  $T_{\text{eff}}$  changes from  $13900 \pm 300$  K, (if obtained from the UVV $\beta$  and uvby $\beta$  photometry), down to  $12850 \pm 50$  K when determined from the Balmer profiles, and can become as low as  $11200 \pm 300$  K when derived from the VI $_c$ J photometry. Furthermore, as far as the Balmer profiles are concerned, the parameters which well reproduce  $H_\beta$ ,  $H_\gamma$ , and  $H_\delta$  predict too narrow  $H_\alpha$  wings. All these discrepancies in the parameter determinations could be accounted for by the presence of the Lupus cloud which could affect the colors in such a way to invalidate the standard reddening relations. Also a spectral contamination by a weak-emission T Tauri (WTT) companion, as suggested by van den Ancker et al. (1996), can not be excluded in an absolute way, although the only spectroscopic sign of its possible presence is a weak variable broad absorption at the position of Li I 6707 Å and a somewhat better agreement between observed and computed spectra longward of 6000 Å for a combined synthetic spectrum obtained from that HR 6000 and one computed for  $T_{\text{eff}}=3500$  K,  $\log g=4.0$ ,  $[M/H]=0.0$ . This companion could be either physically related with HR 6000 to form a close binary system or could be located in the foreground of HR 6000. Finally, the different parameters derived from the different determinations could be due to the extremely peculiar nature of HR 6000 which may make the classical model atmospheres inadequate to reproduce all the observed stellar quantities. In fact, the abundance stratification inferred for some elements, helium in particular, would require more refined models in which the hypothesis of constant abundances throughout the atmosphere is dropped.

By using an ATLAS12 model with parameters  $T_{\text{eff}}=12850$  K,  $\log g=4.1$ ,  $\xi=0.0$  km s<sup>-1</sup> we have computed a synthetic spectrum for HR 6000 from 3050 Å to 9460 Å and have compared it with the observed spectrum. This comparison has shown that most of the elements lighter than Ca are significantly underabundant, except for Be, Na, and P. The Si underabundance ( $[-2.9]$ ) is remarkable because it is even lower than that of 46 Aql ( $[-1.0]$ )


which was claimed by Sadakane et al. (2001) to be the lowest one found in HgMn stars.

A striking peculiarity of HR 6000 is the lack of any overabundance for heavy elements with  $Z > 40$ , except for Xe and Hg. The line spectrum of Xe II is similar to that we observed in a preliminary analysis of UVES spectra of 46 Aql, another Xe rich star: while most of the lines lie at the laboratory wavelength, some other Xe II lines seem to be shifted by about  $-0.1$  Å from the predicted position. The same behaviour can be observed also in HD 175640, although to a less extent, owing to its lower Xe overabundance. Unfortunately, the small number of known transition probabilities and the ignorance of the Xe II isotopic structure due to the lack of atomic data put strong limitations in the study of this element in the spectra of the CP stars.

The large overabundance of  $[+0.7]$  for iron generates a very rich line spectrum in which numerous, still unclassified Fe II lines are observed. A very large number of lines probably due to Fe II remain unidentified. The similar iron overabundance of 46 Aql ( $[+0.65]$ ) gives rise to an impressive close resemblance between the spectra of the two stars.

Abundances in the ultraviolet obtained from a re-analysis of the IUE spectra studied by Castelli et al. (1985) agree with abundances derived from the UVES spectra for all the elements, except for carbon and phosphorus. While the carbon identification in the optical spectrum is rather questionable owing to the weakness of the observed lines, the difference in the phosphorus abundance is similar to that yielded by the individual P II and P III lines.

There are several signs of vertical abundance stratification in HR 6000 for He, P, Mn, and Fe. The peculiar shape of the He I profiles is discussed for the first time in this paper. The profiles have cores too intense as compared to the wings, a fact indicating strong vertical He abundance stratification (Dworetzky, 2004; Bohlender, 2005). Furthermore, the inferred He abundance de-



**Fig. 7.** The observed spectrum of HR 6000 (thick line) is compared with synthetic spectra (thin line) computed with the different choices of Fe II  $\log gf$ 's. The lower panel shows the synthetic spectrum as predicted by the Kurucz (2003) line lists when only transitions from observed levels are considered. Middle panel shows the synthetic spectrum when modified  $\log gf$ 's for several Fe II lines are adopted (Table 6, col. 7). Upper panel displays the synthetic spectrum computed by considering Fe II transitions from both observed and predicted levels.

creases with increasing wavelength ranging from an underabundance of  $[-0.8]$  at  $4000 \text{ \AA}$  to  $[-1.6]$  at  $6000 \text{ \AA}$ .


For Mn II, the abundance shortward of the Balmer discontinuity is larger by about 0.6 dex than that longward of the Balmer discontinuity; for phosphorus, the P II abundance is larger by 0.2 dex than the P III abundance, but this discrepancy lies within the mean square error of the average abundances; for iron, the abundance from Fe II high excitation lines is generally higher by about 0.2 dex than that from Fe II low excitation lines.

Similar to other studied HgMn stars, also HR 6000 shows emission lines. The most numerous ones belong to Mn II, followed by those of Fe II and Cr II. While the first two elements are overabundant in HR 6000, chromium has solar abundance. The star exhibits also isotopic anomalies for Hg and Ca. In both cases the most heavy isotope is the predominant one.

*Acknowledgements.* We are very grateful to J. F. González for the re-reduction of UVES spectra using IRAF routines. and to Dr. J.V. Smoker for the useful discussion regarding interstellar lines.

## References

- Alvarez, E., Arnesen, A., Bengtson, A., Hallin, R., Mayige, C., Nordling, T., Noreland, T., Staaf, O., 1979, *Physica Scripta* 20, 141
- Anders, E., & Grevesse, N. 1989, *Geochim. Cosmochim. Acta*, 53, 197
- Andersen, J., & Jäschek, M. 1984, *A&AS*, 55, 469
- Andersen, J., Jäschek, M., & Cowley, C. R. 1984, *A&A*, 132, 354
- Ballester, P., Grosbol, P., Banse, K., et al. 2000, *SPIE*, 4010, 246
- Bertout, C. 1989, *ARA&A*, 27, 351
- Bessell, M. S., Castelli, F., & Plez, B. 1998, *A&A*, 333, 231
- Bessell, M. S., & Eggen, O. J. 1972, *ApJ*, 177, 209
- Bohlander, D. 2005, *EAS Publications Series*, 17, 83
- Bonifacio, P. 1989, "NORMA: A Program for the Normalization of Spectra", University of Trieste, Astronomy Dept., Internal Rep.20-Apr-1989
- Bonifacio, P., Castelli, F., & Hack, M. 1995, *A&AS*, 110, 441
- Castelli, F. 1999, *A&A*, 346, 564
- Castelli, F., Cornachin, M., Morossi, C., & Hack, M. 1984, *A&A*, 141, 223
- Castelli, F., Cornachin, M., Morossi, C., & Hack, M. 1985, *A&AS*, 59, 1
- Castelli, F., & Hubrig, S. 2004a, *A&A*, 421, L1
- Castelli, F., & Hubrig, S. 2004b, *A&A*, 425, 263
- Castelli, F., & Kurucz, R. L. 1994, *A&A*, 281, 817
- Castelli, F., & Kurucz, R. L. 2004, *ArXiv Astrophysics e-prints*, arXiv:astro-ph/0405087
- Castelli, F., & Kurucz, R. L. 2006, *A&A*, 454, 333
- Catanzaro, G., Leone, F., & Dall, T. H. 2004, *A&A*, 425, 641



**Fig. 8.** Observed and computed profiles in the Li I region. Lower panel shows the spectra observed on September 2005 (thick line) and on March 2006 (dashed line) overplotted on the synthetic spectrum of HR 6000 considered as a single star (thin line). Upper panel compares the spectrum observed on September 2005 (thick line) with the spectrum computed by combining the synthetic spectrum of HR 6000 with a spectrum computed for  $T_{\text{eff}}=3500$  K,  $\log g=4.0$ ,  $[M/H]=0.0$ . The thin labels indicate atomic lines of HR 6000 and the thick labels those of the other star. No labels for molecular lines are printed, although molecular lines were considered for computing the synthetic spectrum of the cool star.

- Smith, K. C. 1997, A&A, 319, 928  
Thé, P. S., Perez, M. R., Voshchinnikov, N. V., & van den Ancker, M. E. 1996, A&A, 314, 233  
Thé, P. S., & Tjin A Djie, H. R. E. 1978, A&A, 62, 439  
van den Ancker, M. E., de Winter, D., & Thé, P. S. 1996, A&A, 313, 517  
Wahlgren, G. M., & Hubrig, S. 2000, A&A, 362, L13  
Wiese, W.L., & Martin, G.A. 1980, NSRDS-NBS, 68  
Zinnecker, H., & Preibisch, T. 1994, A&A, A&A. 292, 152

- Cómerón, F., Fernández, M., Baraffe, I., Neuhäuser, R., & Kaas, A. A. 2003, A&A, 406, 1001
- Cowley, C. R., Hubrig, S., Castelli, F., Gonzalez, J. F., & Wolff, B. 2007, ArXiv Astrophysics e-prints, arXiv:astro-ph/0703227
- Crawford, I. A. 2000, MNRAS, 317, 996
- Crawford, D. L., & Mandwewala, N. 1976, PASP, 88, 917
- Dean, J. F., Warren, P. R., & Cousins, A. W. J. 1978, MNRAS, 183, 569
- Dworetsky, M. M. 2004, IAU Symposium, 224, 727
- Grevesse, N., & Sauval, A. J. 1998, Space Science Reviews, 85, 161
- Hansen, J. E., Persson, W. 1987, Phys. Scr, 36, 602
- Hauck, B., & Mermilliod, M. 1998, A&AS, 129, 431
- James, D. J., Melo, C., Santos, N. C., & Bouvier, J. 2006, A&A, 446, 971
- Johansson, S. 1978, Phys. Scr, 18, 217
- Krautter, J., Wichmann, R., Schmitt, J. H. M. M., Alcalá, J. M., Neuhauser, R., & Terranegra, L. 1997, A&AS, 123, 329
- Kurtz, D. W., & Marang, F. 1995, Delta Scuti Star Newsletter, 8, 2
- Kurucz, R. 1993, SYNTHE Spectrum Synthesis Programs and Line Data. Kurucz CD-ROM No. 18. Cambridge, Mass.: Smithsonian Astrophysical Observatory, 1993., 18,
- Kurucz, R. L. 1997, The Third Conference on Faint Blue Stars, 33
- Kurucz, R. L. 2003, <http://kurucz.harvard.edu/atoms/2601/gf2601.lines0600>
- Kurucz, R. L. 2005, Memorie della Societa Astronomica Italiana Supplement, 8, 86
- Mathis, J. S. 1990, ARA&A, 28, 37
- McWilliam, A. 1998, AJ, 115, 1640
- Moon, T. T. 1985, Commun.Univ.London Obs., 78
- Raassen, A. J. J., & Uylings, P. H. M. 1998, A&A, 340, 300
- Ryabchikova, T. A., Smirnov, Yu. M. 1989, Astron. Tsirk. N1534, 21
- Sadakane, K., et al. 2001, PASJ, 53, 1223
- Siebenmorgen, R., Prusti, T., Natta, A., Müller, T. G. 2000, A&A, 361, 258
- Sigut, T. A. A. 2001, ApJ, 546, L115